


INDONESIAN FISHERIES RESEARCH JOURNAL



AGENCY FOR MARINE AND FISHERIES RESEARCH
MINISTRY OF MARINE AFFAIRS AND FISHERIES

Ind. Fish Res. J.	Vol. 18	No. 2	Page 57-106	December 2012	ISSN 0853-8980
-------------------	---------	-------	-------------	------------------	-------------------

INDONESIAN FISHERIES RESEARCH JOURNAL

Volume 18 Number 2 Desember 2012
Acreditation Number: 503/AU2/P2MI-LIPI/10/2012
(Period: October 2012-October 2015)

Indonesian Fisheries Research Journal is the English version of fisheries research journal.
The first edition was published in 1994 with once a year in 1994. Since 2005, this journal
has been published twice a year on JUNE and DECEMBER.

Head of Editor Board:

Prof. Ir. Badrudin, M.Sc.

Members of Editor Board:

Prof. Dr. Ir. Subhat Nurhakim, M.S.
Prof. Dr. Ir. Ngurah Nyoman Wiadnyana, DEA
Prof. Dr. Ir. Hari Eko Irianto
Dr. Purwito Martosubroto
Dr. Ir. Dede Irving Hartoto, M.Sc.

Refrees for this Number:

Prof. Dr. Ir. Ari Purbayanto, M.Sc.
Prof. Dr. Ir. Setyo Budi Susilo, M.Sc.
Dr. Ir. Augy Syahailatua
Dr. Sonny Koeshendrajana
Dr. Sudarto
Dr. Estu Nugroho

Managing Editors:

Eko Prianto, S.Pi., M.Si.
Arief Gunawan, S.Kom.

Graphic Design:

Kharisma Citra Partadinata, S.Sn.

Published by:

Agency for Marine and Fisheries Research

Manuscript send to the publisher:

Indonesian Fisheries Research Journal
Research Center for Fisheries Management and Conservation
Jl. Pasir Putih I Ancol Timur Jakarta 14430 Indonesia
Phone (021) 64711940; Fax.: (021) 6402640
Email: dewanredaksi@yahoo.com, drprpt2009@gmail.com

Indonesian Fisheries Research Journal is printed by Research Center for Fisheries Management and Conservation Budgeting F.Y. 2012.

PREFACE

Indonesian Fisheries Research Journal Volume 18 Number 2 December 2012 is the second publication of English journal of the Research Center for Fisheries Management and Conservation in 2012. The journal is expected to be a source of newest science and technology for all scientists and researchers in Indonesia and other countries. The financial for publication is provided by the Research Center for Fisheries Management and Conservation budget in the fiscal year of 2012.

This volume contain the Stomach Content of Three Tuna Species in The Eastern Indian Ocean, Exploitation and CPUE Trend of The Small Pelagic Fisheries in The Sulawesi Sea, Indonesia, Is There Any Relationship Between Fluctuating Asymmetry and Reproductive Investment in Giant Featherback (*Chitala lopis*, Notopteridae), Performance of A Fishery Harvesting Different Minimum Shrimp Sizes in The Arafura Sea, Stock Enhancement in Indonesian Lakes and Reservoirs Fisheries, Size and Fishing Ground of Wahoo (*Acanthocybium solandri* Cuvier, 1832) from Catch Data of Tuna Longline Operated in Indian Ocean.

We hope that all the articles on this volume may contribute significantly to the development of fishery science and technology in Indonesia. We are grateful to the editorial board for their improvement and suggestion on reviews of the manuscripts.

Editor

INDONESIAN FISHERIES RESEARCH JOURNAL
Volume 18 Nomor 2 December 2012

CONTENS

Halaman

PREFACE	i
CONTENTS	ii
Stomach Content of Three Tuna Species in the Eastern Indian Ocean <i>By: Bram Setyadji, Andi Bahtiar, and Dian Novianto.....</i>	57-62
Exploitation and CPUE Trend of the Small Pelagic Fisheries in the Sulawesi Sea, Indonesia <i>By: Lilis Sadiyah, Purwanto, and Andhika Prima Prasetyo.....</i>	63-69
Is There any Relationship Between Fluctuating Asymmetry and Reproductive Investment in Giant Featherback (<i>Chitala lopis</i> , Notopteridae) <i>By: Arif Wibowo.....</i>	71-77
Performance of A Fishery Harvesting Different Minimum Shrimp Sizes in the Arafura Sea <i>By: Purwanto.....</i>	79-89
Stock Enhancement in Indonesian Lakes and Reservoirs Fisheries <i>By: Endi Setiadi Kartamihardja.....</i>	91-100
Size and Fishing Ground of Wahoo (<i>Acanthocybium solandri</i> Cuvier, 1832) from Catch Data of Tuna Longline Operated in Indian Ocean <i>By: Agustinus Anung Widodo, Fayakun Satria, and Budi Nugraha</i>	101-106

SIZE AND FISHING GROUND OF WAHOO (*Acanthocybium solandri* Cuvier, 1832) FROM CATCH DATA OF TUNA LONGLINE OPERATED IN INDIAN OCEAN

¹⁾Agustinus Anung Widodo, ²⁾Fayakun Satria and ³⁾Budi Nugraha

¹⁾ Research Center for Fisheries Management and Conservation

²⁾ Research Institute for Fish Resources Rehabilitation and Conservation

³⁾ Research Station for Tuna Fisheries

Received February 2-2012; Received in revised form December 4-2012; Accepted December 5-2012

E-mail: anungwd@yahoo.co.id

ABSTRACT

Wahoo (*Acanthocybium solandri* Cuvier, 1832) is a member of the Scombridae family, is a pelagic (open ocean) species found worldwide in tropical and warm-temperate seas. It is fished throughout its range by artisanal, recreational, and commercial. Wahoo is one of the by-product species of the tuna long line fleets operate in Indian Ocean. This paper describes status of wahoo resource caught by tuna long line in Indian Ocean based at Benoa-Bali. Data obtained from onboard observer program on the tuna long liner based at Benoa-Bali during 2005-2010. Total of 85 trips of onboard observation were carried out with the total long line sets (one set per day) were 2873 times. The data covered the horizontal and vertical position of tuna long line hooks caught the wahoo, hook rate and fish size distribution. Data of horizontal fishing positions (coordinates) gained from the global positioning system available in the tuna long liners. The depth of the long line gear in the waters and temperature of waters were measured by mini-loggers TDR type SP2T-1200, brand: NKE Micrel. Hook rate of wahoo is calculated using the Klawe (1986) method. Result of research showed that the wahoos caught by tuna long lines based at Benoa spread horizontally between 1°31'-33° 40'S and 77°18'-117°53'E and spread vertically between the depth of 75.2- 285.7 m. From 85 tuna long line fishing trips, only about 50% of 85 tuna long line fishing trips caught wahoo with hook rate ranged 0.947-1.399 per 1000 hooks/setting. Size distribution of wahoo ranged 70-180cm with modus ranged 101-110cm.

Keyword: Wahoo, tuna long line, indian ocean

INTRODUCTION

Wahoo (*Acanthocybium solandri* Cuvier, 1832) is known as Pacific kingfish, tiger fish, ocean barracuda, Malata kingfish, queenfish, kingfish. Fishermen in Hawaii called as *ono* and *peto* by fishermen in Karibia. Indonesian fishermen in general called wahoo as *tengiri*, and tuna long line fishermen in Benoa-Bali called as *nyunglas*. Collette & Nauen (1983) described the short description of wahoo as follows, number of dorsal spines (D.XXII-XXVII.12-16), no anal spines, anal soft rays (A.12-14), vertebrae 62 - 64. Mouth large with strong, triangular, compressed and finely serrate teeth. Snout about as long as the rest of head. Posterior part of maxilla completely concealed under pre-orbital bone. Gill rakers absent. Inter-pelvic process small and bifid. Swim bladder present. Body covered with small scales. No anterior corselet developed. The back is iridescent bluish green; the sides silvery with 24 to 30 cobalt blue vertical bars which extend to below the lateral line.

Wahoo (*Acanthocybium solandri* Cuvier, 1832), a member of the Scombridae family, a pelagic (open ocean) species found worldwide in tropical and warm-

temperate seas. The wahoo is highly migratory species, widely distributed throughout tropical and subtropical oceans (Collette & Nauen, 1983; Garber et al., 2005; Mc Bride et al., 2008). Roullot & Venkatasami, (1986) mentioned that sports fishermen used troll around FAD reported good catches of tuna, dolfinfish and wahoo. The major non-tuna catch of what could be considered as the three small-scale fisheries in the region targeting tuna (i.e. those in Cape Verde, Morocco and South Africa) show great differences between the countries. Mostly wahoo in Cape Verde mostly Sarda in Morocco, and almost no non-tuna catch by the small-scale tuna fishery in South Africa (Gillet, 2011). In Indonesia wahoo primarily harvested as bycatch in the troll line, drift gillnet, and tuna long line and purse seine fisheries. This paper describes status of wahoo resource caught by tuna long line in Indian Ocean based at Benoa-Bali.

MATERIALS AND METHODS

Data analyzed were obtained from *onboard observer program* on commercial tuna long line fleets based at Benoa-Bali in the period August 2005 to October 2010. Mostly tuna longlines technology

Corresponding author:

Research Center for Fisheries Management and Conservation Jakarta
Jl. Pasir Putih I Ancol Timur, Jakarta Utara

based at Benoa are using monofilament line that was first introduced into Indonesia by Taiwanese in 1990s. The principal parts of tuna long line consist of a continuous monofilament main line, with floats and branch lines commonly made of smaller gauge monofilament. Numbers of branch line (hook) in between floats were 12 with interval of each branch lines were 50 m. The lengths of each branch lines are 25-30 meter. The deepest hooks usually reach 400 m with target species yellowfin tuna (*Thunnus albacares*) and bigeye tuna (*Thunnus obesus*), whereas wahoo (*Acanthocybium solandri*) was one species caught as by-product. Total of 85 trips were carried out with the total long line sets (one set per day) Of 2873 times. The tuna long line equipped with

radio beacons, radar reflectors, and strobe lights are more likely to be used in marking the gear.

The depth of the long line gear in the waters and temperature of waters were measured by 6 miniloggers TDR type SP2T-1200, brand: NKE Micrel. The minilogger provide information on depth and water temperature, capable to record water depth up to 1200 m and the water temperature -5 to 35°C. The miniloggers attached in the end of branch lines (closed to the hooks). The setting configuration of minilogger was attached to the 1st, 2nd, 3rd, 4th, 5th and 6th branch lines respectively on each basket (Figure 1). With this minilogger setting configuration, each depth range would be covered and recorded.




Figure 1. The setting configuration of the minilogger on the branch line of tuna long line during research.

Identification fish species refer to Collete & Nauen (1983), Compagno (1999), Sainsbury *et al.* (1985) and Sommer *et al.* (1996). The hook rate was calculated by using the equation of Klawe (1986) as follows:

$$HR = \frac{JI}{JP} \times A \quad \dots\dots\dots (1)$$

where: HR = hook rate.
JI = number of fish caught.
JP = number of hook sets.
A = 1000.

RESULTS

Horizotal and Vertival Distribution

Horizontal distribution of wahoo can be described base on result of *onboard observer program* in 2005-2010. The wahoo caught by tuna long line based at

Benoa was between 1°31' S -33° 40'S and 77°18'E - 117°53'E. In 2005-2006, tuna long line based at Benoa-Bali caught wahoo on 12° S -16°S and 107° E -116°E. In 2007, wahoo caught by tuna long line on the area 11° S -34°S and 97° E -115°E. In 2008, area where wahoos were caught 1° S -15°S and 76° E -116° E, whereas in 2009 wahoo caught on the area 1°S-14°S and 96°E-117°E. In 2010 wahoo caught in area 1° S -13°S and 96°E-120°E (Figure 2).

The information on vertical distribution of wahoo was identified through the depth of hook which caught the wahoo. The longline fishing experiment appears as a relevant methodology to achieve this goal (Bach *et al.*, 2003). Installation the *minilogger* on the branch line of tuna long line has been widely used to obtain the information of vertical distribution as used in this research. Result of research show that the wahoo caught by tuna long line spreads between the depths ranged 75- 286 m and mostly caught at depths ranged 100-150 m (Figure 3).




Figure 2. Position of wahoo caught by tuna long line during onboard observer program (2005-2010) which also describes the potential of horizontal distribution of Wahoo.




Figure 3. Depth range that caught by tuna long line during onboard observer program (2005-2010) which also describes the potential of vertical distribution of Wahoo.

Wahoo is a pelagic, highly migratory species of the family scombridae which broadly distributed throughout tropical and subtropical oceans (Collette and Nauen 1983; Garber et al. 2005; Mc Bride et al., 2008). Little information is available on wahoo movement, although their availability change seasonally and the average size different in the various latitude. Wahoo may also migrate seasonally away from the equator following oceanic temperature patterns (Iverson & Yoshida, 1957).

Previous study by Nakano et al. (1997) noted that the catch rate of wahoo in the long line fishery varies at different depth and further argued that wahoo was surface oriented i.e. < 160 meters depth. It is reinforced by evidence the most of wahoo caught by long lines predominantly in the surface (Nakano et al. 1997; Beverly et al., 2009). Sepulveda et al. (2011) noted that depths layer of wahoo is up to 253 m, but in this study recorded that the maximum depth layer of wahoo recorded up to 285.7 m. NMFS (2001) noted

that the distribution of wahoo is associate with warm, productive oceanic fronts, especially when these fronts are near coastal shelves and banks.

Hook Rate

From 85 tuna long line fishing trips, only 42 fishing trips those caught wahoo with total number of catch of wahoo was 463 fishes. Hook rate ranged 0.947-1.399 per 1.000 hooks/setting (Table1, Figure 4). The hook rate is higher compared to the hook rate of wahoo on the tuna long line operated in Pacific Ocean as reported by Kim *et al.* (2007) that the hook rate on the tuna long line operated in Pacific Ocean was 0.446 per 1.000 hooks. The other previous research noted that only 12 wahoo were caught during 5,500 hours of surface trolling in the open sea, more than 60 miles from coastal (Murphy & Ikehara 1955: 14). This species is known to travel solitarily or in small schools of two to twenty fish (Iverson and Yoshida 1957), and is attracted to banks, pinnacles and islands, as well as flotsam drifting in the open ocean (NMFS 2001). That condition allegedly as one of the reasons the hook rate of wahoo on the tuna long line was very low.

Table 1. Hook rate of wahoo on the tuna long line year 2006-2010 based on onboard observer program.

Year	Number of Hooks	Catch of Wahoo (Fish)	Hook Rate (Fish/1000 hooks)
2005	14145	15	1.060
2006	25854	25	0.966
2007	121877	122	1.001
2008	150746	211	1.399
2009	72823	69	0.947
2010	20769	21	1.011




Figure 4. Hook rates fluktuatun of wahoo on the tuna long line fisheries based at Benoa in 2006-2010.

Fish Size

The data of size (fork length-FL) of wahoo caught tuna long line tabulated by year in Table 2. In 2006, the size ranged 72-180 cm, in 2007 ranged 65-145 cm, in 2008 ranged 71-163, in 2009 ranged 70-156cm and in 2010 ranged 109-140 cm. The overall (unsexed) shows that the modus of the size of wahoo was 101-110 cm as delineated on Figure 5.

Table 2. Size of wahoo caught tuna long line based in Benoa during year 2006-2010.

Years	Number of sample (fish)	Fork length (cm)	
		Min	Max
2006	21	72	180
2007	17	65	145
2008	33	71	163
2009	48	70	156
2010	18	109	140




Figure 5. Size distribution of wahoo caught tuna long line based at Benoa in 2006-2010.

Sommer *et al.* (1996) noted that maximum total length (TL) of wahoo was 250 cm (unsexed), length at first maturity L_m ranged 85-105 cm and the modus at 99.3 cm and according to Collette (1995) the common fork length was 170 cm (unsexed). Refer to this fact, the wahoo caught by tuna long line in Indian Ocean based in Benoa mostly were adult fish. This evidence is also indicates the tuna long line was selective fishing gear for wahoo.

CONCLUSION

From above description can be concluded as follows, the wahoo caught by tuna long line based at Benoa was between 1°31' S -33° 40'S and 77°18'E - 117°53'E and spreads between the depth of 75.2-285.7 m. The hook rate ranged 0.947-1.399 per 1000 hooks/setting and mostly were adult fish.

ACKNOWLEDGMENTS

We express our thank to observers of Research Institute for Tuna Fishery Benoa Mr. Abram Barat, Mr. Dian Novianto, Mr. Andi Bachtiar, Mr. Gamadi, Mr. Yusuf Affandi, Mr. I Nyoman S., Mr. Irwan Jatmiko as well as Mr. I Gede Sanjaya who provided the data. We also thank to Director of RCMFC Purwanto, PhD for valuable inputs and comments.

REFERENCES

- Bach, P., Dagorn, L., Bertrand, A., Josse, E., & Misselis, C. 2003. Acoustic telemetry versus monitored longline fishing for studying the vertical distribution of pelagic fish: bigeye tuna (*Thunnus obesus*) in French Polynesia. *Fisheries research*, 60: 281-292.
- Beverly, S., Curran D., Musyl M., & Molonya B. 2009. Effects of eliminating shallow hooks from tuna longline sets on target and non-target species in the Hawaii-based pelagic tuna fishery. *Fish Res* 96:281-288.
- Kim, S.S., D.H. An, D.Y. Moon, & S.J. Hwang, 2007. Comparison of circle hook and J hook catch rate for target and bycatch species taken in the Korean tuna long line fishery during 2005-2006. *Scientific Committee Third Regular Session*. 13-24 August 2007, Honolulu, United of America. WCPFC-SC3-EB SWG/WP-11. 11 p.
- Poisson, F. 2006. Compilation of information on neritic tuna species in the Indian Ocean. A Working Paper, IOTC Secretariat. IOTC-2006-SC-INF11. *IOTC Ninth Session of the Scientific Committee*, Mahé, Seychelles, 6-10 November 2006. 23 p.
- Roullot, J. & A. Venkatasami. 1986. Dispositifs de Concentration de Poissons (DCP): l'expérience mauricienne. *Doc. FAO*. TWS/86/26. *Expert consultation on the stock assessment of tunas in the Indian Ocean*. Colombo, 4-8 December 1986. 12 p.
- Garber A.F., Tringali M.D., & Franks J.S., 2005. Population genetic and phylogeographic structure of wahoo, *Acanthocybium solandri*, from the western central Atlantic and central Pacific Oceans. *Mar Biol*. 147: 205-214.
- Gillett, R. 2011. Bycatch in small-scale tuna fisheries. A global study. *FAO Fisheries And Aquaculture Technical Paper* No.560. Food And Agriculture Organization Of The United Nations, Rome. 116 p.
- Collette, B.B. 1995 Scombridae. Atunes, bacoretas, bonitos, caballas, estorninos, melva, etc. In W. Fischer, F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter and V. Niem (eds.) *Guía FAO para Identificación de Especies para lo Fines de la Pesca. Pacífico Centro-Oriental*. 3 Vols. FAO, Rome. p. 1521-1543.
- Collette, B.B. and C.E. Nauen. 1983. *FAO Species Catalogue*. Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. *FAO Fish. Synop*. Rome: FAO. 125 (2): 137 p.
- Sommer, C., W. Schneider and J.-M. Poutiers 1996. *FAO species identification field guide for fishery purposes. The living marine resources of Somalia*. FAO, Rome. 376 p.
- Iverson, E. & H. Yoshida. 1957. *Notes on the biology of the wahoo in the Line Islands*. Pacific Science. 11: 370-379.
- Nakano, H., M. Okazaki & O. H. 1997. Analysis of catch depth by species for tuna longline fishery based on catch by branch lines. *Bull Nat Res Inst Far Seas Fish*. 34: 43-62.
- NMFS. 2001. Final Environmental Impact Statement: Pelagic Fisheries of the Western Pacific Region. Prepared by: URS Corporation, 615 Piikoi St, Ste 900, Honolulu HI 96814. Southwest Fisheries Science Center, NMFS/NOAA. Available online: <http://swr.nmfs.noaa.gov/piao/eisdocs.htm>.
- Stevens, M.M. 2004. Wahoo/Ono, *Acanthocybium solandri*. Wahoo_Pacific_SFW_MMS_Final Report.doc. *Seafood Watch-Seafood Report*. Monterey Bay Aquarium.
- Mc Bride R.S., Richardson A.K., Maki K.L. 2008. Age, growth, and Mortality of wahoo, *Acanthocybium solandri*, from the Atlantic coast of Florida and the Bahamas. *Mar Fresh Res*. 59:799-807.

Sepulveda C.A., Scott A. Aalbers, Soûa Ortega Garcia, Nicholas C.Wegner & Diego Bernal. 2011. Depth distribution and temperature preferences of wahoo (*Acanthocybium solandri*) off Baja California Sur, Mexico. *Mar Biol.* 158:917–926.

INDONESIAN FISHERIES RESEARCH JOURNAL

Instructions for Authors

Scope

Indonesian Fisheries Research Journal publishes research results on resources, oceanography and limnology for fisheries, fisheries biology, management, socio-economic and enhancement, resource utilization, aquaculture, post harvest, of marine, coastal and inland waters.

Manuscripts submission

Manuscripts submitted to the Journal must be original with clear definition of objective, materials used and methods applied and should not have been published or offered for publication elsewhere. The manuscripts should be written in English, in double-spaced typing on one side A4-size white papers. **Three copies** of the manuscripts and a disk containing the manuscript file (s) produced using MS Word must be submitted to the **Managing Editor, Agency for Marine and Fisheries Research, Jl. Pasir Putih II, East Ancol, Jakarta 14430, Indonesia, Tlp. (021) 64700928, Fax. (021) 64700929. E-mail: drprpt2009@gmail.com. Website: <http://p4ksi.litbang.kkp.go.id/>**. There are no page charges for the manuscripts accepted for publications, unless otherwise indicated below.

Manuscripts preparation

1. A **title page** should be provided which shows the title and the name(s) of the author(s). The author(s)' position and affiliation should be written as footnotes at the bottom of the title page.
2. **Title** should not be more than 15 words.
3. **Abstract** summarizes the study is not more than 250 words. Keywords (3-5) must be provided and conform with Agrovocs.
4. **Introduction** should concisely establish the relate to which the article concerns, the purpose and the importance of the work. Do not use any sub-headings.
5. **Materials** and **Methods** should clearly and concisely describe the experiment with sufficient detail for independent repetition.
6. **Results** are presented with optimum clarity with unnecessary detail. No same results should be presented in both tables and figures. **Tables** (double spaced, same font size with text, no vertical line, lower case supercript letters to indicate, footnotes), and **illustration/figures** should be printed on separate sheets and given arabic number consecutively. **illustration/figures** should be submitted in original (1) and copied (2) forms and be sufficiently sharp for reproduction. 1 **Photographs** are preferably black and white on glossy papers with good contrast. Color photographs of good quality are acceptable but the cost of reproduction must be paid for by the author(s). Exemption for bearing the cost can be sought from the publisher by written request.
7. **Conclusion** should be short with regard to the title, objectives, and discussion of results.
8. **Acknowledgement**, if necessary, should be kept minimum (less than 40 words).
9. **References** should be cited in the text by the author(s)' family or last name and date in one or two forms: Wang (1985) or (Wang, 1985). For references with more than two authors, cite the first author plus *et al.* Full citation in alphabetical order is required for the references list in the following style:

Sunarno, M.T.D., A. Wibowo & Subagja. 2007. Identifikasi tiga kelompok ikan belida (*Chitala lopis*) di Sungai Tulang Bawang, Kampar dan Kapuas dengan pendekatan biometrik. *Jurnal Penelitian Perikanan Indonesia*. 13 (3): 1-14.

Sadhotomo, B. 2006. Review of environmental features of the Java Sea. *Indonesia Fisheries Research Journal*. 12 (2): 129-157.

Collins, A. 1977. Process in acquiring knowledge. In Anderson, R.C., Spiro, R.J. and Montaque, W.E. (eds.). *Schooling and the Acquisition of Knowledge*. Lawrence Erlbaum, Hillsdale, New Jersey. p.339-363

Bose, A.N., Ghosh, S.N., Yang, C.T. and Mitra, A. 1991. *Coastal Aquaculture Engineering*. Oxford & IBH Pub. Co. Pvt. Ltd., New Delhi. 365 pp.

Smith, T.I.J. and Sandifer, P.O. 1983. Development of prawn (*Macrobrachium rosenbergii*) farms in temperate climates: Prospects and problems in the United States. In Roger, G.L., Day, D. and Um, A. (eds.) *Proceedings of the First International Conference on Warm Water Aquaculture - Crustacea*. Brigham Young University, Laie, Hawaii, USA. p 109-126.

Unpublished materials should not be used, except for thesis which is cited as follows:

Simpson, B.K. 1984. *Isolation, Characterization and Some Applications of Trypsin from Greenland Cod (Gadus morhua)*. PhD Thesis. Memorial University of New Foundland, St. John's, New Foundland, Canada. 179 pp.

10. **Acronyms or uncommon abbreviations** must be given in full with the first mentions; new abbreviations should be coined only for unwieldy name and should not be used at all unless the names occur frequently.
11. **Latin name** and **family of the species** should be given besides its common name at first mention in the manuscript, and the common name only for the next mentions.
12. **SI Systems** should be used for all measurements.
13. **Reprint**: A free reprint of the journal issue in which the paper is published will be sent to the author.



**LEMBAGA
ILMU PENGETAHUAN
INDONESIA**



**Panitia
Penilai
Majalah
Ilmiah**

SERTIFIKAT

Nomor: 503/AU2/P2MI-LIPI/10/2012

Akreditasi Majalah Ilmiah

Kutipan Keputusan Kepala Lembaga Ilmu Pengetahuan Indonesia
Nomor 893/E/2012 Tanggal 1 Oktober 2012

Nama Majalah : Indonesian Fisheries Research Journal
ISSN : 0853-8980
Redaksi : Pusat Riset Perikanan Tangkap
Kementerian Kelautan dan Perikanan

Ditetapkan sebagai Majalah Ilmiah

TERAKREDITASI

Akreditasi sebagaimana tersebut di atas berlaku selama 3 (tiga) tahun

Cibinong, 1 Oktober 2012

Lembaga Ilmu Pengetahuan Indonesia
Ketua Panitia Penilai Majalah Ilmiah-LIPI

Prof. Dr. Rochadi

NIP 195007281978031001



Certified company
Cert No.: 13.001.1773



KAN
Kantor Akreditasi Nasional
Lembaga Sertifikasi Sistem Manu
LSM 016 IDN

