

THE ASSESSMENT OF SEDIMENT CONTAMINATION LEVEL IN THE LAMPUNG BAY, INDONESIA: HEAVY METAL PERSPECTIVE

Fitri Budiyanto¹⁾ & Lestari¹⁾

¹⁾Research Center for Oceanography, Indonesian Institute of Sciences

Diterima tanggal: 23 Februari 2015; Diterima setelah perbaikan: 5 Mei 2015; Disetujui terbit tanggal 1 Juli 2015

ABSTRAK

Teluk Lampung memiliki nilai sosial-ekonomi dan ekologi yang tinggi berkaitan dengan potensi perairan dan penggunaannya oleh manusia. Di lain pihak, pemanfaatan Teluk Lampung mungkin mengubah kelimpahan bahan kimia berbahaya seperti logam berat. Tujuan dari penelitian ini untuk mengetahui konsentrasi logam berat Cd, Cu, Pb dan Zn dalam sedimen dan menilai kondisi perairan Teluk Lampung. Pengamatan konsentrasi logam berat dilakukan di 13 titik stasiun pada bulan maret 2008. Analisis logam berat dalam sedimen menggunakan tiga jenis asam: HNO₃, HCl, H₂O₂ kemudian sampel dianalisis menggunakan Spektrofotometer Serapan Atom. Hasil pengukuran menunjukkan variasi logam berat di setiap lokasi pengamatan dan konsentrasi tertinggi Cd, Cu, Pb dan Zn dalam sedimen secara berurutan adalah 0.08 mg/kg berat kering, 22.99 mg/kg berat kering, 24.75 mg/kg berat kering dan 118.48 mg/kg berat kering. Faktor yang dominan mempengaruhi variasi logam berat dalam sedimen dalam studi ini adalah jarak lokasi pengamatan dengan pusat kegiatan antropogenik dan fraksi sedimen. Indeks SQG-Q menunjukkan 7 titik stasiun memiliki nilai SQG-Q ≤ 0.1 sementara 6 titik stasiun lainnya memiliki nilai 0.1 ≤ SQG-Q < 1. Hal ini berarti lebih dari setengah titik stasiun berada pada kondisi tidak tercemar.

Kata kunci: Teluk Lampung, konsentrasi logam berat, indeks SQG-Q

ABSTRACT

The potency and utilization of Lampung Bay has been recognized for their socio-economical and ecological values. However, heavy use of this Bay may alter the abundance of hazardous chemical like heavy metals. The aims of this study were to determine the concentration of Cd, Cu, Pb and Zn in the sediment and to assess Lampung Bay water condition. The observation of heavy metal content in sediment of Lampung Bay was conducted at 13 stations in March 2008. Analysis of heavy metals in sediment was conducted using three kinds of acid: HNO₃, HCl and H₂O₂ while measurement was carried out by Atomic Absorption Spectrophotometer. The result indicated a variation of heavy metal concentration in sediment and that concentrations of Cd, Cu, Pb and Zn in sediment were 0.08 mg/kg dry weight, 22.99 mg/kg dry weight, 24.75 mg/kg dry weight and 118.48 mg/kg dry weight, respectively. Factors influenced heavy metal concentration in sediment in this study including the distance between sampling location and anthropogenic activities and the sediment fraction. SQG-Q index indicated that 7 stations have SQG-Q ≤ 0.1 whereas other 6 stations have 0.1 ≤ SQG-Q < 1, meaning that more than half sampling stations are in uncontaminated state.

Keywords: Lampung Bay, heavy metals concentration, SQG-Q index

INTRODUCTION

Heavy metals enrich water body by several mechanisms (Lee & Morel, 1985) and their abundance in sediment can represent level of contamination of the environment (Lestari & Witasari, 2010). Riverine runoff becomes major source of metals in estuarine waters (Darmono, 1995) and depositional processes alter composition of metal in the sediment (Rompas, 2010). Changes of physicochemical properties (pH, salinity, temperature, eH) provoke the release of metals in sediment into upper water column (Hosono *et. al.*, 2010). Some of the released metals were redeposited in the sediment and this process will be repeated several times inflicting permanent release of metals into water column and/or permanent sink of metals into seabed sediment (Rejomon *et. al.*, 2010).

Lampung bay is located in the southernmost part of Sumatra that shows socio-economical and ecological value due to its marine resources and utilization (Fitriya, 2010; Thoha, 2010). Mangrove, coral reef, seagrass,

sand and mud are part of ecosystem diversity in Lampung Bay (Cappenberg, 2010; Handayani, 2010). Coastal communities utilize those resources for aquaculture activities, such as: pearl farming, seaweed aquaculture and tourism (Kadi, 2010; Pramudji, 2010).

Local Government of Lampung (1999) explained that Lampung receives spill-over of economic growth of Jakarta causing rapid enhancement of anthropogenic activities like industry, urban area, agriculture and aquaculture. According to Susana *et. al.*, (2001), Lampung bay has been influenced by industrial activities like cement, coal, timber and oil production. These activities may alter metals concentration in aquatic ecosystem especially Cd, Cu, Pb and Zn which will be used and will be released in their production process (Darmono, 1995). The aims of this research were to determine Cd, Cu, Pb and Zn concentration in sediment and to assess its contamination level.

Korespondensi Penulis:

Jl. Pasir Putih I Ancol Timur, Jakarta Utara 14430. Email: fitr019@lipi.go.id

METHODOLOGY

Sampling site

Sediment samples were collected in 13 observation stations in March 2008 in the geographical area of 105.22° to 105.37° East and 5.66° to 5.46° South (Figure 1). The stations were chosen representing local anthropogenic activities. Northern part of the Bay represents urban areas, and the western part is proximal to aquaculture area. The eastern part of the bay was selected for observation in the water near industrial area.

The collection of sediment samples

In each station, sample was taken using stainless steel grab sampler then 0-10 cm of surface-layer sediment was collected using Polyethylene spoon. In the field, sediment sample was stored in ice box at ± 4°C in temperature (Hutagalung *et. al.*, 1997).

Analytical method

Acidic destruction and spectrophotometry measurement were used to analyze sediment samples according to USEPA 3050b procedure (USEPA, 1996). HNO₃ (1+1) was added into a gram of dry sediment then the solution was heated for 15 minutes. After the heating, concentrated HNO₃, H₂O₂ 30% and concentrated HCl were added by dropper into solution and reheating of solution was performed. Aquadest was added into sample solution to 100 ml in volume. After the destruction processes finished, Cd, Cu, Pb

and Zn in sediment were measured using Flame Atomic Absorption Spectrophotometer (FAAS) Varian SpectrAA 20 plus utilizing air-acetylene as ignition gas. Before used, all glasswares were soaked into HNO₃ (1+1) for 24 hours and were rinsed by aquadest.

To understand Cd, Cu, Pb and Zn concentration in sediment, sediment fraction was measured then PCA analysis was conducted to discover the influence of sediment fraction to the metals concentration. Statistica version 6.0 was used to perform PCA analysis and the sedimentary fraction data was taken from Helfinalis (2010) as a secondary data.

Index Calculation

Ecosystem changes by anthropogenic activities in this research were evaluated using SQG-Q (mean sediment quality guideline quotient) index. SQG-Q was used to determine pollution level in the bed sediment by comparing with quality guideline. Caeiro *et. al.*, (2005) formulated SQG-Q index as:

$$SQG - Q = \frac{\sum_{i=1}^n PEL-Q_i}{n} \dots\dots\dots 1)$$

$$PEL - Qi = \frac{contaminant}{PEL} \dots\dots\dots 2)$$

- PEL-Qi : probable effect level quotient for each contaminant
- PEL : probable effect level for each contaminant (Tabel 1)
- Contaminant : metals concentration in sediment
- n : number of parameter for calculation




Figure 1. Sampling stations for sediment sample in the Lampung Bay.

Table 1. The references for SQG-Q index

Element	Cd	Cu	Pb	Zn
Probable effect level (PEL) value for each metal, mg/kg dry weight	4,2	108	112	271

Source:Nobi *et. al.*, 2010

Table 2. Interpretation of SQG-Q value

SQG-Q value	Designation of sediment quality
SQG-Q ≤ 0,1	Unimpacted: lowest potential for observing adverse biological effects
0,1 < SQG-Q ≤ 1	Moderate impact potential for observing adverse biological effects
SQG-Q ≥ 1	Highly impacted potential for observing adverse biological effects

In this calculation, PEL values were used to figure out the index value (Table 1). The others value could be used to substitute PEL like quality standard of metals in the sediment, even though, substituting this value possibly changes the calculated index.

The value obtained from those calculations varied from 0 up to 1. Caeiro *et. al.*, (2005) explained the meaning of the value of SQG-Q and the brief of that interpretation was summarized in Table 2.

RESULT AND DISCUSSION

Metals concentration varied among locations, and both anthropogenic activities and natural processes alter metals abundance. Cd became the lowest value and the tightest range of metals found in the bed sediment compared with Cu, Pb and Zn. On the opposite, Zn had the widest range of value among the others (Table 3). Cd shows the worst toxic effect and its presence in sediment probably affects benthic organism even in low concentration (Ellwood, 2004). Cd, Cu and Pb naturally exist in oil and coal then the utilizing of both oil and coal

as fuel may release the metals into aquatic ecosystems. Zn has been categorized as essential metal which is required and is used in metabolism, however, recent researches showed that excessive uptake of Zn into organism body gives toxic effect (Dean *et. al.*, 2007). Report of Zn hazards deccribed that Zn deficiency have severe effects on all stages of growth, development, reproduction and survival, however, exceeding level of zinc causes negative effect.

Urban and aquaculture activities along Lampung Bay coast line influenced metals abundance. The high concentration of Cu, Pb and Zn were detected in the river mouth and urban area, and those metals tend to decrease by the distance. Cd evenly distributed in this Lampung Bay, for Cd naturally presents in a little amount in earth crust (Taylor, 1964), so little changes in enviroment have no significant effect to Cd concentration. Sanusi (2006) explained that river flow includes human waste such as metals contaminant. Other researchers reported that increasing metals content in the river mouth was affected by input volume/runoff of river containing metals (Rochyaton *et.*

Table 3. Concentration of Cd, Cu, Pb and Zn in Sediment

Station	Metal in sediment concentration mg/kg dry weight			
	Cd	Cu	Pb	Zn
st 1	0.05	20.58	21.61	111.00
st 2	0.05	22.99	24.73	118.48
st 3	0.08	18.27	6.05	99.06
st 4	0.03	5.30	7.28	35.27
st 5	0.04	16.88	12.20	71.65
st 6	0.04	12.84	11.94	61.67
st 7	0.01	11.01	10.01	63.79
st 8	0.02	8.39	9.23	33.14
st 9	0.03	8.82	9.22	57.52
st 10	0.03	9.06	9.29	63.11
st 11	0.04	8.33	9.49	58.62
st 12	0.02	7.12	9.88	52.99
st 13	0.02	8.35	9.99	54.01

al., 2005; Ciutat et. al., 2007). Rochyatun et. al., (2006) observed the impact of riverine runoff in the Cisadane river as long as the fluctuation of metals content on bed sediment by flooding phenomenon. Their research discovered that Cd and Zn increased in two fold of concentration after the flooding, however, Pb and Cu showed no differences between pra and post of the flooding. Urban activities contribute in metals input mostly by fuel combustion, paint weathering, metal furniture corrosion and urban waste disposal (Ciutat et. al., 2007). Aquaculture area contributes in Cu and Zn input into aquatic ecosystem. Dean et. al., (2007) reported the highest Cu and Zn concentration found in fish farming area which feed and faecal of fish contain of Cu and Zn. In addition, Cu is used as antifouling in boat paint (Olsgard, 1999).

Sediment fraction also affected Cd, Cu, Pb and Zn concentration in bed sediment where clay fraction

strongly bound Cd, Cu, Pb and Zn (Figure 2). Clay and silt concentrated in the urban and aquaculture area (Table 5) where Cd, Cu, Pb and Zn found in relatively high concentration. Considering the industrial area in the eastern part of the Bay have coarser sediment than the urban area metals concentration is in low concentration. Finer particles provide larger surface area, and these kinds of particle bind more metals onto particle. Moreover, finer particles form dense bed sediment inhibiting the release of trapped-metals into water column (Donazzolo et. al., 1984; Martinctic et. al., 1990). River mouth and urban area of Lampung Bay tend to contain more clay and resulted in high concentration of metals detected in those areas (Figure 2).

Metals concentration in bed sediment of Lampung Bay is considered as natural concentration of metals in sediment. Cd, Cu, Pb and Zn concentration in sediment

Table 4. Concentration of Cd, Cu, Pb and Zn in sediment in the other locations

Location	Metals concentration in sediment, mg/kg dry weight				Reference
	Cd	Cu	Pb	Zn	
East Kalimantan, Indonesia	<0.001-0.1	2.02-14.5	4.4-15.2	15.8-121.2	Rochyatun, et. al., 2003
Pari Islands, Indonesia	0.03-0.06	13.24-15.93	0.84-6.01	0.73-17.83	Rochyatun, 2003
Semarang, Indonesia	0.06-0.13	18.3-36.6	10.9-17.3	13.6-16.3	Lestari, 2011
Brest Harbour, France	-	700	500	1160	Cabon, et. al., 2010
Masan Bay, Korea	1.24	43.40	43.97	-	Hyun, et. al., 2007
Mexican Caribbean	-	1.3	1	-	Whelan III, et. al., 2011
Andaman Islands, India (sandy ecosystem)	0.69-1.96	6.64-7.04	-	10.4-27.72	Nobi, et. al., 2010
Andaman Islands, India (dead coral ecosystem)	1.24-1.75	5.48-10.58	-	25.54-38.77	Nobi, et. al., 2010
Andaman Islands, India (seagrass ecosystem)	1.04-3.88	44.36-86.76	3.08-6.64	21.64-48.72	Nobi, et. al., 2010
Andaman Islands, India (mangrove ecosystem)	0.8-1.52	80.86-87.93	3.9-5.4	12.21-23.02	Nobi, et. al., 2010
Abundance in the continent crust	0.2	55	12.5	70	Taylor, 1964
Lampung bay, Indonesia	0.01-0.08	5.30-22.99	6.05-24.73	33.14-118.48	This study

Table 5. Grain Size classification of the sediment in each station of The Lampung Bay

Station	Sediment Classification (%) ^a					Remark
	Pebbles	Granules	Sand	Silt	Clay	
st 1	0	0	4,32	52,64	43,04	Urban area
st 2	0	1,83	5,74	27,5	64,93	Urban area
st 3	0	0	14,22	42,2	43,57	River Mouth
st 4	0	4	35,45	22,04	38,51	-
st 5	0	0	1,66	48,87	50,46	Aquaculture area
st 6	0	0	6,02	53,41	40,57	Aquaculture area
st 7	9,14	8,52	11,78	28,55	42,02	Aquaculture area
st 8	13,73	11,67	12,96	23,48	38,16	Industrial area
st 9	0	0,38	13,55	56,6	29,47	Aquaculture area
st 10	0	7,16	18,68	33,78	40,39	Aquaculture area
st 11	0	0,73	3,55	32,34	63,37	Industrial area
st 12	0	0,74	4,36	54,63	40,27	Industrial area
st 13	0	0	3,92	47,05	49,03	-




Figure 2. Principal Components Analysis for sediment fraction and concentration of metals in sediment.

Table 6. SQG-Q index for each station

Station	PEL-Qi				SQG-Q
	Pb	Cd	Cu	Zn	
st.1	0,193	0,0114	0,191	0,410	0,20
st.2	0,221	0,0124	0,213	0,437	0,22
st.3	0,054	0,0188	0,169	0,366	0,15
st.4	0,065	0,0074	0,049	0,130	0,06
st.5	0,109	0,0100	0,156	0,264	0,13
st.6	0,107	0,0100	0,119	0,228	0,12
st.7	0,089	0,0026	0,102	0,235	0,11
st.8	0,082	0,0038	0,078	0,122	0,07
st.9	0,082	0,0062	0,082	0,212	0,10
st.10	0,083	0,0076	0,084	0,233	0,10
st.11	0,085	0,0088	0,077	0,216	0,10
st.12	0,088	0,0050	0,066	0,196	0,09
st.13	0,089	0,0050	0,077	0,199	0,09

of Lampung Bay showed no extreme differences with the average abundance within continental crust in preindustrial period. Metal concentration in sediment in the Lampung Bay was lower than other areas like harbor and other urban waters (Table 4). This study supported the report in metal's bioaccumulation in Lampung Bay (Tugiyono, 2007) that discovered that Hg and Pb concentration in Lampung Bay was not dangerous for marine organisms. The concentration of Hg and Pb in green mussle (*Perna viridis*) increased, yet those concentrations surpassed none of existing regulation.

SQG-Q index

SQG-Q index value varied among station as the metals concentration differed among locations. Zn contributed the most in the calculation of SQG-Q index than other metals due to the high concentration of Zn compared with PEL value. Interpreting the SQG-Q index, 7 stations (more than a half of total stations) were unimpacted area and moderate impact was found in the rest of the stations (Table 6). The presence of metals in sediment will expose metals accumulation in benthic organism in various mechanisms. The effect of metals accumulation in organism differs in each metal, depending on: intake rate, exposure route, internal

factor of organisms (ages, sex) and physiochemical properties of environment (Luoma & Rainbow, 2008). Cd and Pb categorized as non-essential element having toxic effect even in low concentration. In contrast, Cu and Zn were essential elements used in metabolisms, however, exceeding concentration has toxic potential. In recent research, Cd was found to substitute Zn and its function in carbonic anhydrase enzyme of diatomae when the seawater lacked Zn. In this state, diatomae could maintain its growth rate, however, this phenomenon still is not be observed in higher trophic level organisms (Lane *et. al.*, 2005).

SQG-Q index calculates the sum of measured concentration and quality guideline (in this study, PEL was used), and it predicts toxicity of metals in sediment. The achieved score predicted the influence of metal's abundance in benthic organism, however, real adverse effect of metals to organism could not be determined by these scores only. The effect of metals bioaccumulation to marine organism is complex (Luoma & Raibow, 2008). The changes of water properties may affect the toxicity of sediment. The presence of other chemical components like sulphide or organic material influence metals bioavailability. In metal-sulphide compound, metals were strongly bound and in this state metals mobility considered low, resulted in low toxicity effect of sediment to marine organism. Furthermore, the existence of other metals caused different impact on toxicity, particularly interaction of metal with one another. In most cases, the poisoning effect of Cd can be prevented by Zn consumption and the consumption of Cd decreases Cu concentration in the liver (Darmono, 1995).

CONCLUSION

Anthropogenic activities along Lampung Bay coastal area altered Cd, Cu, Pb and Zn concentration in bottom sediment. Cu, Pb and Zn concentrated in urban and aquaculture area which represent human activity. However, Cd distributed evenly in the Bay due to low concentration of Cd in the natural sediment indicating another factor, such as sediment fraction also contributed in its concentration. Urban and aquaculture area are covered by finer sediment than other areas in the Lampung Bay resulted in high metals concentration. Comparison with previous study areas, Cd, Cu, Pb and Zn in the Lampung Bay were relatively low. Calculation of SQG-Q index showed that more than half of the total observation stations were claimed to unimpacted area and the others were in moderate impact state.

ACKNOWLEDGEMENTS

This study was funded by DIPA Research Centre for Oceanography-LIPI. We acknowledge Prof. Pramudji as researcher in chief and our deep thanks to

Mr. Abdul Rozak and Mr. M. Taufik Kaisupy for assisting laboratory analysis.

REFERENCES

- Cabon, J.Y., P. Giamarchi & S.L. Floch. (2010). A study of marine pollution caused by the release of metals into seawater following acid spills. *Marine Pollution Bulletin* 60: 998-1004.
- Caeiro, S., M.H. Costa, T.B. Ramos, F. Fernandes, N. Silveira, A. Coimbra, G. Medeiros & M. Painho, (2005). Assessing heavy metal contamination in Sado Estuary sediment: An index analysis approach. *Ecological Indicators* 5: 151-169.
- Cappenberg, H.A.W., (2010). Komposisi jenis dan distribusi moluska di perairan Teluk Lampung. In: Ruyitno dkk. (Ed.). *Status Sumber daya laut di perairan Teluk Lampung*: 14-24. Jakarta: LIPI Press.
- Ciutat, A., M. Gerino, & A. Boudou, (2007). Remobilization and bioavailability of cadmium from historical contaminated sediments: Influence of bioturbation by tubificids. *Ecotoxicology and Environmental Safety* 68: 108-117.
- Darmono. (1995). *Logam dalam sistem biologi makhluk hidup*. Jakarta: UI Press. 136 hlm.
- Dean, R.J., T.M. Shimmield & K.D. Black, (2007). Copper, zinc and cadmium in marine cage fish farm sediments: An extensive survey. *Environmental Pollution* 145: 84-95.
- Donazzolo, R., O. Hieke-Merlin, L. Menagazzo-Vitturi & B. Pavoni, (1984). Heavy metal content and lithological properties of recent sediments in the Northern Adriatic. *Marine Pollution Bulletin* 15: 93-101.
- Ellwood, M.J., (2004). Zinc & cadmium speciation in subantarctic waters east of New Zealand. *Marine Chemistry* 87: 37-58.
- Fitriya, N., (2010). Kelimpahan zooplankton di perairan Teluk Lampung. In: Ruyitno dkk. (Ed.). *Status Sumber daya laut di perairan Teluk Lampung*: 104-112. Jakarta: LIPI Press.
- Handayani, T., (2010). Makroalgae ekonomis penting di perairan Teluk Lampung. In: Ruyitno dkk. (Ed.). *Status Sumber daya laut di perairan Teluk Lampung*: 25-31. Jakarta: LIPI Press.
- Helfinalis, (2010). Lingkungan pantai, endapan sedimen dan total suspended solid di Perairan

- Teluk Lampung. In: Ruyitno dkk. (Ed.). Status Sumber daya laut di perairan Teluk Lampung: 205-219. Jakarta: LIPI Press.
- Hosono, T., C.C. Su, F. Siringan, A. Amano & S.I. Onodera, (2010). Effect of environmental regulations on heavy metal pollution decline marine core sediments from Manila Bay. *Marine Pollution Bulletin* 60: 780-785.
- Hutagalung, H.P., Setiapermana, D. & Riyono, S.H., (1997). Metode analisis air laut, sedimen dan biota, Buku 2. Jakarta: P3O LIPI. 182 hlm.
- Hyun, S., C.H. Lee, T. Lee & J.W. Choi, (2007). Anthropogenic contributions to heavy metal distributions in the surface sediments of Masan Bay, Korea. *Marine Pollution Bulletin* 54: 1031-1071.
- Kadi, A., (2010). Karakteristik komunitas rumput laut beserta interaksi antar jenis di perairan Teluk Lampung. In: Ruyitno dkk. (Ed.). Status Sumber daya laut di perairan Teluk Lampung: 51-62. Jakarta: LIPI Press.
- Lane, T.W., M.A. Saito, G.N. George, I.J. Pickering, R.C. Prince, & F.M.M. Morel, (2005). A cadmium enzyme from a marine diatom. *Nature Publishing Group* 435: 42.
- Lee, J.G. & F.M.M. Morel. (1985). Replacement of zinc by cadmium in marine phytoplankton. *Marine Ecology Progress Series*. 127: 305-309.
- Lestari, (2011). Distribusi dan geokimia logam berat dalam sedimen di Perairan Semarang, Jawa Tengah. In: A. Hartoko dkk. (Ed.). *Prosiding Pertemuan Ilmiah Nasional Tahunan VIII ISOI 2011*: 204-217. Makassar: Ikatan Sarjana Oseanologi Indonesia.
- Lestari & Y. Witasari. (2010). Kualitas perairan di Teluk Lampung ditinjau dari aspek logam berat, In: Ruyitno dkk. (Ed.). Status Sumber daya laut di perairan Teluk Lampung: 160-170. Jakarta: LIPI Press.
- Local Government of Lampung, (1999). Atlas sumberdaya wilayah pesisir lampung. Bandar Lampung: Pemerintah Daerah Provinsi Lampung. 80 hlm.
- Luoma, S.N. & P.S. Rainbow, (2008). Metal contamination in aquatic environments: Science and lateral Management. United Kingdon: Cambridge University press. 556 hlm.
- Martincic, D., Z. Kwokal & M. Branica, (1990). Distribution of zinc, cadmium and copper between different size fractions of sediments I. The Limski Kanal (North Adriatic Sea). *The Science of the Total Environment* 95: 201-215.
- Nobi, E.P., E. Dilipan, T. Thangaradjou, K. Sivakumar & L. Kannan, (2010). Geochemical and geo-statistical assessment of heavy metal concentration in the sediments of different coastal ecosystems of Adaman Islands, India. *Estuarine, Coastal and Shelf Science* 87: 253-264.
- Olsgard, F., (1999). Effect of copper contamination on recolonisation of subtidal marine soft sediment- an experimental field study. *Marine Pollution Bulletin* 38: 448-462.
- Pramudji, (2010). Mangrove di kawasan pesisir Teluk Lampung, Provinsi Lampung. In: Ruyitno dkk. (Ed.). Status Sumber daya laut di perairan Teluk Lampung: 1-13. Jakarta: LIPI Press.
- Rejomon, G., M. Nair & T. Joseph, (2010). Trace metal dynamics infishes from the Southwest coast of India. *Environment Monitoring Assessment*. 167: 243-255.
- Rochyatun, E., Edward & A. Rozak, (2003). Kandungan logam berat Pb, Cd, Cu, Zn, Ni, Cr, Mn & Fe dalam air laut dan sedimen di perairan Kalimantan Timur. *Oseanologi dan Limnologi di Indonesia* 35: 51-71.
- Rochyatun, E., (2003). Kadar logam berat dalam air laut dan sedimen di Perairan Gugus Pulau Pari. In: Ruyitno dkk. (Ed.). *Pesisir dan Pantai Indonesia VIII*: 99-104. Jakarta: P2O-LIPI.
- Rochyatun, E., Lestari & A. Rozak, (2004). Kondisi perairan Muara Sungai Digul dan Perairan Laut Arafura dilihat dari kandungan logam berat. *Oseanologi dan Limnologi di Indonesia*. 36: 15-31.
- Rochyatun, E., Lestari & A. Rozak, (2005). Kualitas lingkungan perairan Banten dan sekitarnya ditinjau dari kondisi logam berat. *Oseanologi dan Limnologi di Indonesia*. 38: 23-46.
- Rochyatun, E., M.T. Kaisupy & A. Rozak. (2006). Distribusi logam berta dalam air dan sedimen di perairan muara Sungai Cisadane. *Makara Sains*. 10 (1): 35-40.
- Rompas, R.M., (2010). Toksikologi kelautan. Jakarta: Sekretariat Dewan Kelautan Indonesia. 75 hlm.
- Sanusi, H.S., (2006). Kimia Laut: Proses fisik kimia

dan interaksinya dengan lingkungan. Bogor: Departemen Ilmu dan Teknologi Kelautan, Fakultas Perikanan dan Ilmu Kelautan, Institut Pertanian Bogor. 188 hlm.

Susana, T, Suyarso & M. Muchtar, (2001). Karakteristik beberapa parameter kimia, kaitannya dengan tataguna lahan perairan Teluk Lampung. In: Aziz, A and M. Muchtar (Ed.). Perairan Indonesia: Oseanografi, Biologi dan Lingkungan: 43-53. Jakarta: P3O LIPI.

Taylor, S.R., (1964). Abundance of chemical elements in the continental crust: a new table. *Geochimica et Cosmochimica Acta*.28: 1273-1285.

Thoha, H., (2010). Fitoplankton di perairan Teluk Lampung. In: Ruyitno dkk. (Ed.). Status Sumber daya laut di perairan Teluk Lampung: 96-103. Jakarta: LIPI Press.

Tugiyono. (2007). Bioakumulasi logam Hg dan Pb di perairan Teluk Lampung, Propinsi Lampung. *Jurnal Sains MIPA*. 13 (1): 44-48.

United State of Environmental Protection Agency, (1996). Test methods for evaluating solid waste. SW-846 Methods 3050B.

Whelan III, T., B.I. Van Tussenbroek & M.G.B. Santos, (2011). Changes in trace metals in *Thalassia testudinum* after hurricane impact. *Marine Pollution Bulletin* 62: 2797-2802.